

Heavy Lifting Causing Spontaneous Coronary Artery Dissection

with Anterior Myocardial Infarction in a 54-Year-Old Woman

Kyriakos Yiangou, MD Kyriacos Papadopoulos, MD Chara Azina, MD Spontaneous coronary artery dissection in association with strenuous exercise and weightlifting is rather sparsely described in the medical literature. Diagnosis and treatment of this rare condition is a challenge, but prompt recognition and appropriate early choice of angioplasty or surgery can lead to a good outcome.

We report the case of a postmenopausal 54-year-old woman who presented with anterior myocardial infarction caused by spontaneous dissection of the left anterior descending coronary artery after she had lifted a heavy weight while gardening. The patient was treated successfully by means of angioplasty and the implantation of 3 drug-eluting stents. In addition to presenting the patient's case, we review the topical medical literature. **(Tex Heart Inst J 2016;43(2):189-91)**

pontaneous coronary artery dissection (SCAD), a rare cause of myocardial infarction (MI) and sudden cardiac death, can present with various acute coronary symptoms. It usually occurs in the peripartum period. In association with strenuous exercise and heavy lifting, SCAD is rather sparsely documented in the medical literature. We report the case of a patient in whom heavy lifting caused SCAD, and we review the literature on this topic.

Case Report

In January 2014, a 54-year-old woman had acute-onset retrosternal chest pain immediately after gardening and disposing of a heavy basket of cut grass. She had a normal body weight, no relevant medical history, and no known risk factors for coronary artery disease. She was 4 years postmenopausal. After an hour of worsening pain, she presented at the emergency department, having experienced cardiac arrest and resuscitation en route. After she was intubated, an electrocardiogram (ECG) showed ST-segment elevation in leads V₂ through V₅, so she underwent thrombolysis. The patient's troponin T level was elevated at 0.4 ng/mL. The next day, her ECG results were consistent with anterior MI. Coronary angiograms revealed a spiral dissection of the left anterior descending coronary artery (LAD) (Fig. 1), which was successfully treated by means of angioplasty and the implantation of 3 drug-eluting stents. The rest of the coronary tree was smooth. The patient was discharged from the hospital after an uneventful 6-day stay, with instructions to take aspirin, prasugrel, a β -blocker, and a statin. Seven months later, results of an exercise ECG were normal, and the patient had returned to normal activity and full-time work.

Discussion

Coronary artery dissection refers to the splitting of the arterial wall layers, the result of which is a false lumen. This dissection can occur spontaneously; after chest trauma, percutaneous coronary intervention, or cardiac surgery; or as part of aortic dissection. Accumulated blood in the false lumen can encroach on the true lumen, thereby impairing blood flow and causing myocardial ischemia, MI, or sudden cardiac death.¹⁻⁵ Since the first description of SCAD,⁶ approximately 320 documented cases have been reported. This figure might be misleading, because many of the reports were

Key words: Aneurysm, dissecting/complications/ diagnosis/etiology/radiography; myocardial infarction/

etiology; risk factors; treatment outcome; weight lifting/injuries

From: Cardiology Department (Dr. Yiangou), The Cardio Clinic Heart Centre, 2042 Nicosia; Cardiology Department (Dr. Papadopoulos), Medical Check-up Centre, 2014 Nicosia; and Internal Medicine Department (Dr. Azina), Nicosia General Hospital, 2230 Latsia, Nicosia; Cyprus

Address for reprints:

Kyriakos Yiangou, MD, Leoforos Strovolou 131, 2042 Nicosia, Cyprus

E-mail:

yiangouk@hotmail.com

© 2016 by the Texas Heart® Institute, Houston of autopsy findings.⁷ The overall incidence of SCAD ranges from 0.3% to 1.1%.⁷⁸ Most victims are parturient women.⁵⁹ The LAD is involved in 60% of cases.⁵

Spontaneous coronary artery dissection is often associated with coronary atherosclerosis,^{5,8} wherein the local inflammatory process in the plaque disrupts the normal adherence of the intima and media, and hence causes flap formation. In autopsy studies, eosinophilic infiltration has been the most likely explanation for dissection during the peripartum period.^{10,11} Eosinophils contain lytic substances, such as peroxidase and collagenase, that break down the medial-adventitial layers, leading to dissection. Increased levels of estrogen and



Fig. 1 Coronary angiograms in *A*) right anterior oblique and *B*) anteroposterior views show the left anterior descending coronary artery dissection (arrows).

progesterone during pregnancy and childbirth might also exaggerate the eosinophilic presence.^{12,13} Hormones and hemodynamic factors might cause microstructural changes in the elastic and collagen fibers of the aorta.

Other clinical conditions have been associated with SCAD, such as hypertension,¹⁴ Marfan syndrome,¹⁵ Ehlers-Danlos syndrome,¹⁶ polyarteritis nodosa,¹⁷ systemic lupus erythematosus,¹⁸ eosinophilic arteritis,¹⁹ and rarer forms of vasculitis that involve the arterial wall. Medications such as oral contraceptives,²⁰ fenfluramine,²¹ and cyclosporine²² have been implicated. Coronary spasm might worsen variant angina,²³ and cocaine use²⁴ has also been reported as a cause of SCAD.

Among the cardiovascular sequelae of lifting heavy weight, the chief culprit is vascular stress on arterial walls: during maximal exertion, blood pressure can reach 480/350 mmHg.²⁵ This mechanical compression and associated phenomena might have caused the SCAD in our patient, who was postmenopausal and had no known coronary risk factors.

The diagnosis of SCAD is usually made using coronary angiography. Also, intravascular ultrasound,²⁶ multislice computed tomography,²⁷ and optical coherence tomography²⁸ yield images of excellent quality.

Treatment for SCAD involves medical therapy and interventional procedures or surgery. The ultimate choice depends upon the individual clinical and angiographic characteristics of each patient. Because so few cases have been reported, there are no guidelines to suggest an appropriate management algorithm. Although thrombolysis has been used with success in patients who have SCAD, primary angioplasty has been recommended as the chief treatment.²⁹ We used both in our patient's case.

Spontaneous coronary artery dissection is a rare yet well-recognized cause of acute coronary syndromes. In addition to weightlifting,³⁰ certain predisposing factors and conditions should raise clinical suspicion of SCAD. Prompt diagnosis and appropriate management with early angioplasty are important to optimize clinical care and ensure a good outcome.

References

- 1. Auer J, Punzengruber C, Berent R, Weber T, Lamm G, Hartl P, Eber B. Spontaneous coronary artery dissection involving the left main stem: assessment by intravascular ultrasound. Heart 2004;90(7):e39.
- Bergen E, Huffer L, Peele M. Survival after spontaneous coronary artery dissection presenting with ventricular fibrillation arrest. J Invasive Cardiol 2005;17(10):E4-6.
- Kurum T, Aktoz M. Spontaneous coronary artery dissection after heavy lifting in a 25-year-old man with coronary risk factors. J Cardiovasc Med (Hagerstown) 2006;7(1):68-70.
- El-Sherief K, Rashidian A, Srikanth S. Spontaneous coronary artery dissection after intense weightlifting UCSF Fresno Department of Cardiology. Catheter Cardiovasc Interv 2011;78 (2):223-7.

- Kamran M, Guptan A, Bogal M. Spontaneous coronary artery dissection: case series and review. J Invasive Cardiol 2008; 20(10):553-9.
- 6. Pretty HC. Dissecting aneurysm of coronary artery in a woman aged 42: rupture. Br Med J 1931;1(3667):667.
- Vanzetto G, Berger-Coz E, Barone-Rochette G, Chavanon O, Bouvaist H, Hacini R, et al. Prevalence, therapeutic management and medium-term prognosis of spontaneous coronary artery dissection: results from a database of 11,605 patients. Eur J Cardiothorac Surg 2009;35(2):250-4.
- Hering D, Piper C, Hohmann C, Schultheiss HP, Horstkotte D. Prospective study of the incidence, pathogenesis and therapy of spontaneous, by coronary angiography diagnosed coronary artery dissection [in German]. Z Kardiol 1998;87(12):961-70.
- 9. Thompson EA, Ferraris S, Gress T, Ferraris V. Gender differences and predictors of mortality in spontaneous coronary artery dissection: a review of reported cases. J Invasive Cardiol 2005;17(1):59-61.
- Koul AK, Hollander G, Moskovits N, Frankel R, Herrera L, Shani J. Coronary artery dissection during pregnancy and the postpartum period: two case reports and review of literature. Catheter Cardiovasc Interv 2001;52(1):88-94.
- Robinowitz M, Virmani R, McAllister HA Jr. Spontaneous coronary artery dissection and eosinophilic inflammation: a cause and effect relationship? Am J Med 1982;72(6):923-8.
- Tchernitchin AN, Barrera J, Arroyo P, Mena MA, Vilches K, Grunert G. Degranulatory action of estradiol on blood eosinophil leukocytes in vivo and in vitro. Agents Actions 1985;17 (1):60-6.
- Grunert G, Tchernitchin AN. Effect of progesterone on the non-genomic response to oestrogen in the rat. J Endocrinol 1982;94(3):307-15.
- Basso C, Morgagni GL, Thiene G. Spontaneous coronary artery dissection: a neglected cause of acute myocardial ischaemia and sudden death. Heart 1996;75(5):451-4.
- Bateman AC, Gallagher PJ, Vincenti AC. Sudden death from coronary artery dissection. J Clin Pathol 1995;48(8):781-4.
- Ades LC, Waltham RD, Chiodo AA, Bateman JF. Myocardial infarction resulting from coronary artery dissection in an adolescent with Ehlers-Danlos syndrome type IV due to a type III collagen mutation. Br Heart J 1995;74(2):112-6.
- Chu KH, Menapace FJ, Blankenship JC, Hausch R, Harrington T. Polyarteritis nodosa presenting as acute myocardial infarction with coronary dissection. Cathet Cardiovasc Diagn 1998;44(3):320-4.
- Aldoboni AH, Hamza EA, Majdi K, Ngibzadhe M, Palasaidi S, Moayed DA. Spontaneous dissection of coronary artery treated by primary stenting as the first presentation of systemic lupus erythematosus. J Invasive Cardiol 2002;14(11):694-6.
- Lepper PM, Koenig W, Moller P, Perner S. A case of sudden cardiac death due to isolated eosinophilic coronary arteritis [published erratum appears in Chest 2005;128(5):3779]. Chest 2005;128(2):1047-50.
- Evangelou D, Letsas KP, Korantzopoulos P, Antonellis I, Sioras E, Kardaras F. Spontaneous coronary artery dissection associated with oral contraceptive use: a case report and review of the literature. Int J Cardiol 2006;112(3):380-2.
- Goli AK, Koduri M, Haddadin T, Henry PD. Spontaneous coronary artery dissection in a woman on fenfluramine. Rev Cardiovasc Med 2007;8(1):41-4.
- 22. Tsimikas S, Giordano FJ, Tarazi RY, Beyer RW. Spontaneous coronary artery dissection in patients with renal transplantation. J Invasive Cardiol 1999;11(5):316-21.
- Mark DB, Kong Y, Whalen RE. Variant angina and spontaneous coronary artery dissection. Am J Cardiol 1985;56(7):485-6.

- 24. Jaffe BD, Broderick TM, Leier CV. Cocaine-induced coronary-artery dissection. N Engl J Med 1994;330(7):510-1.
- MacDougall JD, Tuxen D, Sale DG, Moroz JR, Sutton JR. Arterial blood pressure response to heavy resistance exercise. J Appl Physiol (1985) 1985;58(3):785-90.
- Deftereos S, Giannopoulos G, Mavrogianni A, Sykiotis A, Pyrgakis V, Bobotis G. Role of grey-scale intravascular ultrasound and ChromaFlo in deciding on treatment approach for spontaneous coronary dissection in a young woman. Hellenic J Cardiol 2011;52(4):364-6.
- 27. Ney L, Korner M, Leibig M, Heindl B. Traumatic dissection of a coronary artery: detection by multislice computed tomography and use of tirofiban as a reversible platelet inhibitor. Resuscitation 2011;82(3):358-60.
- Iglesias D, Salinas P, Jimenez-Valero S. Spontaneous coronary artery dissection evaluated by optical coherence tomography. J Cardiovasc Med (Hagerstown) 2011;12(10):743-4.
- 29. Andreou AY, Georgiou PA, Georgiou GM. Spontaneous coronary artery dissection: report of two unsuspected cases initially treated with thrombolysis. Exp Clin Cardiol 2009;14 (4):e89-92.
- Fred HL. More on weightlifting injuries. Tex Heart Inst J 2014;41(4):453-4.