

Vertebral Doppler Bunny Waveform

Suggesting Subclavian Stenosis and Influencing
Coronary Artery Bypass Technique

Sergio Suma, MD
Stefano Coli, MD
Alessandro Palumbo, MD
Stefania Graziuso, MD
Bruno Borrello, MD
Nicola Gaibazzi, MD

A 73-year-old man presented with acute coronary syndrome and 3-vessel coronary disease and underwent carotid ultrasonography before surgical revascularization. His medical history included right carotid stenting and iliofemoral bypass. He had no substantial blood pressure differences between arms, and no obvious subclavian or carotid bruits.

A pulsed-wave Doppler ultrasonogram revealed a “bunny” waveform pattern of the left vertebral artery. This pattern, characterized by a systolic peak, a midsystolic dip, and a more blunted systolic peak (Fig. 1), suggested left subclavian stenosis.¹ Computed tomographic angiograms showed irregular plaque, constituting 60% to 70% stenosis of the left subclavian artery and extending toward the origin of the left vertebral artery (Figs. 2 and 3).

To avoid the effect of upstream subclavian stenosis and possible coronary subclavian steal after revascularization,² we prepared a left internal mammary artery bypass to the left anterior descending coronary artery as a free graft.

Comment

The bunny waveform resembles a rabbit’s profile. It is thought to be caused by post-stenotic pressure decrease resulting from high-velocity turbulent flow through the ste-




Fig. 1 Pulsed-wave Doppler ultrasonogram of the left vertebral artery shows a “bunny” waveform: a midsystolic dip (arrow), a first systolic peak (the rabbit’s ears), and a second blunted systolic peak (the rabbit’s rump), suggesting left subclavian stenosis.



Fig. 2 Reconstructed computed tomographic angiogram shows irregular plaque, constituting 60% to 70% stenosis of the left subclavian artery (arrowhead) and extending toward the origin of the left vertebral artery (arrow).

Section Editor:
Raymond F. Stainback, MD

From: Cardiology Department (Drs. Coli, Gaibazzi, and Suma), Radiology Department (Drs. Graziuso and Palumbo), and Cardiac Surgery Section (Dr. Borrello), Azienda Ospedaliero-Universitaria di Parma, 43126 Parma, Italy

Address for reprints:
Sergio Suma, MD,
Via Golfo Dei Poeti 3,
43126 Parma, Italy

E-mail:
sergiosuma.md@gmail.com

© 2020 by the Texas Heart®
Institute, Houston




Fig. 3 Computed tomographic angiograms (3-dimensional reconstruction) in left **A**) anterolateral and **B**) posterolateral views show irregular plaque, constituting 60% to 70% stenosis of the left subclavian artery (arrowheads) and extending toward the origin of the left vertebral artery (arrows).

nosis during systole. The degree of midsystolic velocity decrease, which is associated with the severity of subclavian stenosis, indicates a “pre-steal” state that precedes systolic flow reversal seen in more advanced disease.

The usefulness of carotid ultrasonography before surgical revascularization is debated.³ Current guidelines rate the need for cerebrovascular duplex examination before cardiac surgery as “uncertain” in all clinical situations, particularly in asymptomatic patients before coronary artery bypass grafting.⁴ More research is needed in this area.

Recognizing the vertebral Doppler bunny waveform pattern in our patient led to the diagnosis of clinically unsuspected subclavian stenosis and to an important change in our surgical approach to revascularization.

References

1. Kliever MA, Hertzberg BS, Kim DH, Bowie JD, Courneya DL, Carroll BA. Vertebral artery Doppler waveform changes indicating subclavian steal physiology [published erratum appears in AJR Am J Roentgenol 2000;174(5):1464]. AJR Am J Roentgenol 2000;174(3):815-9.
2. Sintek M, Coverstone E, Singh J. Coronary subclavian steal syndrome. Curr Opin Cardiol 2014;29(6):506-13.
3. Masabni K, Sabik JF 3rd, Raza S, Carnes T, Koduri H, Idrees JJ, et al. Nonselective carotid artery ultrasound screening in patients undergoing coronary artery bypass grafting: is it necessary? J Thorac Cardiovasc Surg 2016;151(2):402-8.
4. American College of Cardiology Foundation (ACCF); American College of Radiology (ACR); American Institute of Ultrasound in Medicine (AIUM); American Society of Echocardiography (ASE); American Society of Nephrology (ASN); Intersocietal Commission for the Accreditation of Vascular Laboratories (ICAVL), et al. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS/SVU 2012 appropriate use criteria for peripheral vascular ultrasound and physiological testing part I: arterial ultrasound and physiological testing [published erratum appears in J Am Coll Cardiol 2013;62(16):1540]. J Am Coll Cardiol 2012;60(3):242-76.